

## LA-UR-20-22978

Approved for public release; distribution is unlimited.

Title: Heavy Flavor and jet studies for the future Electron Ion Collider

Author(s): Li, Xuan

Intended for: APS April meeting 2020, 2020-04-18/2020-04-21 (Washington, District Of Columbia, United States)

Issued: 2020-04-20 (rev.1)

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

# Heavy Flavor and jet studies for the future Electron Ion Collider

Xuan Li ([xuanli@lanl.gov](mailto:xuanli@lanl.gov))

on behalf of Los Alamos National Laboratory




Xuan Li (LANL)

## Outline


- Motivation.
- Initial detector design and tracking performance.
- Open heavy flavor studies in simulation.
- Summary and outlook.

# New QCD frontier: the Electron Ion Collider (EIC)

- EIC can help solve different fundamental physics problems in a wide  $x$  and  $Q^2$  kinematic region.
  - How quarks and gluons distributed in momentum and space within the nucleon and a heavy nuclei?
  - **Proton spin origin?**
  - What happens to the gluon density in nuclei, does it saturate at high energy?
  - A clean environment to study the flavor dependent energy loss in nuclear medium.



A. Accardi et al, *Eur. Phys. J. A*, 52 9 (2016).






# New EIC heavy flavor and jet program proposed by LANL (I)

- Through measuring heavy flavor hadrons, jets which can be treated as surrogates of initial quarks/gluons and their correlations in the hadron/nuclei going (forward) direction at the EIC.

[arXiv:1610.08536](https://arxiv.org/abs/1610.08536)

Phys. Rev. D 96, 114005 (2017)


$$e^- + p \rightarrow e^- + jet(D^\pm) + X$$



- To precisely determine the initial quark/gluon distribution functions in the poorly constrained kinematic region.
- To precisely study the quark/gluon fragmentation/hadronization processes.
- To provide further information on the gluon Sivers function and other spin observables.

# New EIC heavy flavor and jet program proposed by LANL (II)


- Through measuring heavy flavor hadrons, jets which can be treated as surrogates of initial quarks/gluons and their correlations in the hadron/nuclei going (forward) direction at the EIC.



- To understand the nuclear medium effects on hadron production such as modification on nuclear PDFs, parton energy loss mechanisms and hadronization processes through the comparison of measured hadron/jet cross section between e+p and e+A collisions.

# New EIC physics observables are under study


- Competing models of nuclear modification in DIS reactions with nuclei (e.g HERMES data). Differentiation not possible with light hadrons.
  - Hadronization inside nuclear matter (dashed lines).
  - Energy loss of partons, hadronization outside the nuclear matter (solid lines).
- Heavy mesons have very different fragmentation functions and formation times
  - Easy to discriminate between larger suppression for D/B mesons (in-medium hadronization) and strong/intermediate  $z$  enhancement (E-loss).
  - Enhanced sensitivity to the transport properties of nuclei.
- A Forward Silicon Tracking (FST) detector is needed at the future EIC to carry out such measurements.




Projected stat. uncertainties at event generation level and include evaluated sampling efficiencies.

# Initial design of the proposed Forward Silicon Tracking detector

- Initial detector design in fast simulation:
  - Assumed mid-rapidity silicon vertex detector with 3 barrel layers of Monolithic Active Pixel Sensor (MAPS) type sensors.
  - Forward-rapidity silicon tracking detector (FST): 2 barrel layers of MAPS silicon detector and 4 forward planes of MAPS + other silicon detector.





$B = 3T$   
FST:  $1.0 < \eta < 4.0$



# Tracking performance of the proposed FST

- Evaluated track performance of the FST with pixel pitch  $30\text{ }\mu\text{m}$ , material budgets per detector layer:  $0.3\%X_0$  and the readout rate is at  $500\text{ kHz}$  :




- Better than  $70\text{ }\mu\text{m}$  resolution can be achieved by the initial FST design for the **transverse decay length  $b_T$  measurements** for tracks with  $p_T > 1\text{ GeV/c}$  over the  $1.5 < \eta < 3.5$  region.
- The **momentum resolution  $dp_T/p_T$**  are better than or consistent with the forward tracking requirements from the EIC detector handbook.


# Reconstructed D mesons with the proposed FST in simulation

- The full analysis framework which includes the event generation (PYTHIA8), detector response in fast simulation, beam remnant interaction background embedding, and hadron reconstruction have been setup.
- Mass distributions of clusters with track transverse decay length matching between charged tracks with smeared tracking performance on momentum/spatial resolutions with integrated luminosity:  $10 \text{ fb}^{-1}$ .


Reconstructed cluster mass with  $K^\pm$



Reconstructed cluster mass with  $K^\pm$




Reconstructed cluster mass with  $K^\pm$



- Track  $\eta$ : 0.5 to 4
- Tracking efficiency set at 95%.
- The performances are based on 100%  $K/\pi/p$  separation.


# Flavor dependent nuclear modification factor projections for reconstructed hadrons

- Reconstructed other heavy flavor hadrons with the help of the proposed FST, e.g.  $\Lambda_c^+$ ,  $B^\pm$ .



- The statistical precision of reconstructed hadrons can help separate different models of the nuclear modification on hadronization processes.
- Heavy flavor measurements** at the EIC will enhance the sensitivity of the nuclear transport properties.




## Projected hadron RAA vs $z_h$





# Summary and Outlook

- The new heavy flavor and jet program for the EIC will shed light into the flavor dependent energy loss and parton fragmentation processes in the poorly constrained kinematic region.
- Ongoing detector and physics simulation studies will provide guidance on the detector requirements.
- Ongoing EIC detector R&D and detector design will provide realistic detector performance parameters for the simulation studies.
- We look forward to work with more collaborators and contribute to the EIC realization.










# Backup

# Physics performance on detector reference

- Early simulation work done by JLab colleagues on  $D^0$  meson reconstruction with different detector systems.

[arXiv:1610.08536](https://arxiv.org/abs/1610.08536)

$D^0 \rightarrow \pi K$  mass spectrum, on the top of DIS background



# EIC detector performance requirements


- From the EIC detector handbook.

Table 2: Physics requirements for an EIC detector

EIC Detector Requirements															
$\eta$	Nomenclature			Tracking			Electrons		$\pi/K/p$ PID		HCAL	Muons			
				Resolution	Allowed $X/X_0$	Si-Vertex	Resolution $\sigma_E/E$	PID	p-Range (GeV/c)	Separation	Resolution $\sigma_E/E$				
-6.9 — -5.8	↓ p/A	Auxiliary Detectors	low- $Q^2$ tagger	$\delta\theta/\theta < 1.5\%$ ; $10^{-6} < Q^2 < 10^{-2} \text{ GeV}^2$											
...															
-4.5 — -4.0			Instrumentation to separate charged particles from photons				2%/√E								
-4.0 — -3.5															
-3.5 — -3.0	Central Detector	Backwards Detectors	$\sigma_p/p \sim 0.1\% \times p + 2.0\%$	$\sim 5\%$ or less	TBD	$\pi$ suppression up to 1:10 <sup>4</sup>		$\leq 7 \text{ GeV}/c$	$\geq 3\sigma$	$\sim 50\%/\sqrt{E}$					
-3.0 — -2.5															
-2.5 — -2.0											$\sigma_p/p \sim 0.05\% \times p + 1.0\%$	7%/√E			
-2.0 — -1.5															
-1.5 — -1.0															
-1.0 — -0.5		Barrel	$\sigma_p/p \sim 0.05\% \times p + 0.5\%$		$\sigma_{xyz} \sim 20 \text{ }\mu\text{m}$ , $d_0(z) \sim d_0(r\phi) \sim 20/p_T \text{ GeV }\mu\text{m} + 5 \text{ }\mu\text{m}$		(10-12)%/√E	$\leq 5 \text{ GeV}/c$		$\geq 3\sigma$	TBD	TBD			
-0.5 — 0.0															
0.0 — 0.5															
0.5 — 1.0													Forward Detectors	$\sigma_p/p \sim 0.05\% \times p + 1.0\%$	TBD
1.0 — 1.5															
1.5 — 2.0		$\sigma_p/p \sim 0.1\% \times p + 2.0\%$			$\leq 20 \text{ GeV}/c$										
2.0 — 2.5															
2.5 — 3.0															
3.0 — 3.5															
3.5 — 4.0	↑ e	Auxiliary Detectors	Instrumentation to separate charged particles from photons							$\leq 45 \text{ GeV}/c$					
4.0 — 4.5															
...															
> 6.2					Proton Spectrometer			$\sigma_{\text{intrinsic}}( d )/ d  < 1\%$ ; Acceptance: $0.2 < p_T < 1.2 \text{ GeV}/c$							

# LANL EIC program progress – inclusive jet simulation

- Moreover, the FST enables the flavor tagged jet and jet substructure studies at forward rapidity for the EIC.
- Initial look into the kinematics of inclusive jets in e+p collisions:



PYTHIA8 MC  
 $E_e = 18 \text{ GeV}$   
 $E_p = 100 \text{ GeV}$   
 $0.0 < \eta < 4.5$   
Anti-kT  
algorithm  
 $R = 1.0$   
Int. Luminosity:  
 $10 \text{ fb}^{-1}$

- Jet flavor tagging study is under way.